

# What is the Endocannabinoid System and What is its Role?

---

Terrence Shenfield MS, RRT-ACCS, RPFT, NPS, AE-C





Blue Dream (1/S)	<b>THC</b>
	15.93%
	<b>CBD</b>

Purple Cindy 99 (S/D)	<b>THC</b>
	15.00%
	<b>CBD</b>

Purple	<b>THC</b>
	12.84%

Island Sweet Skunk (S)	<b>THC</b>
	15.50%

Lime	<b>THC</b>
	11.80%
	<b>CBD</b>


Mango Sou Jack (S/M)	<b>THC</b>
	12.50%
	<b>CBD</b>




# Objectives

- History of cannabis in the USA
- The endocannabinoid system
- Endocannabinoid Deficiency
- Side effects and drug interactions
- CB1 and CB2 Receptors





Cannabidiol



Raphael Mechoulam




# Endocannabinoids: *Bind CB1 > CB2* *structure, related to prostaglandins*

- **Anandamide**

(arachidonyl-ethanolamid)

- **2-Arachidonoyl - glycerol (2-AG)**

more abundant, less potent




# **Understanding Endocannabinoid Deficiency**



The cannabinoid receptors are further divided into 2 main subtypes, known as cb1 and cb2.



**CB1** is found mostly in the brain.




**CB2** receptors are mostly found within the immune system.



# Background and History

- Cannabis is thought to have originated in central Asia (probably China).
- There is evidence of hemp rope dating back to 8,000 B.C.







# Background and History in US

- 1600-1890's Hemp production encouraged
- 1906 Pure Food and Drug Act required labeling of hemp products
- 1900-1920's Mexican use of recreational marijuana
- 1930's Fear of marijuana – Linked to violence and deviate behavior
- 1930 Creation of the Federal Bureau of Narcotics (FBN)
  - Harry J. Anslinger was the first Commissioner of the FBN and remained in that post until 1962





# Background and History in US

- 1932 Uniform State Narcotic Act
  - Required States to handle to epidemic
- 1936 Reefer Madness propaganda film
- 1937 Marijuana Tax Act- Congress and the “Evil Weed” for medical and industrial use
- 1944 La Guardia Report finds marijuana less dangerous
- 1940’s- Hemp used for WW2 military equipment





# Background and History in US

- 1951-1956- Stricter penalties for possession
- 1960's Counter culture use of marijuana
  - President Kennedy and President Johnson said doesn't lead to other drugs and violence
- 1968 Creation of the Bureau of Narcotics and Dangerous Drugs
  - This was a merger of FBN and the Bureau of Dangerous Drugs of the Food and Drug Administration.






# Background and History in US

- 1972 Shafer Commission
  - Decriminalize marijuana but president Nixon rejected the report
- 1973 Creation of the DEA
- 1976 Beginning of parents' movement against marijuana
- 1986 president Nixon – 3 strikes and your out equated marijuana with heroin
- 1996 Medical Use in California




# Quick History of the Receptors



# Cannabinoid Receptors


G protein-coupled

- **CB1**
  - Brain, fat cells, liver, duodenum, muscle
- **CB2**
  - Lymphocytes > macrophages > cytokines



# Depolarization-induced suppression of inhibition

**POSTSYNAPTIC** endocannabinoid release inhibits **PRESYNAPTIC** GABA and glutamate release



Nicoll & Alger, 2004



# The Human Endocannabinoid System

The endocannabinoid system consists of two receptors, called CB1 and CB2. These receptors are found on cell surfaces and impact various biological processes.

**CB<sub>1</sub>**


Located in the brain, central nervous system, and many other parts of the body.

**CB<sub>2</sub>**

Found throughout the body on cells associated with our immune system.

## Cannabidiol (CBD)

CBD is one of the primary cannabinoids found in hemp. It interacts with **CB<sub>1</sub>** and **CB<sub>2</sub>** receptors for many effects still being studied.



# Endocannabinoid Overview

- Endogenous physiologic system
  - Establishes and maintains normal state or homeostasis
  - Key role in food hunger, fat accumulation, glucose and lipid metabolism
- Two well known receptor site
  - CB<sub>1</sub> –central and peripheral nervous system
  - CB<sub>2</sub> –immune cells
- Endogenous endocannabinoids
  - AEA (Anandamide)
  - 2-AG (2-Arachidonoylglycerol)
- Enzymes degrade AEA and 2-AG

Sources

<http://noml.org/library/item/introduction-to-the-endocannabinoid-system>

<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2241751/>

These statements have not been evaluated by the FDA and are not intended to diagnose, treat or cure any disease.






# Cannabinoid Receptors:

- CB1
  - THC has high affinity for these receptors
  - Pain , nausea, and depression
  - Arthritis and Lupus
- CB2
  - Receptors part of immune system
  - Great densities found in GI tract
  - Modulate Crohn's disease and IBS



# CB1 physiologic effects

- CB1 blockade helps
  - Regulate energy intake
  - Improves glucose and lipid metabolism
  - Antinociception
  - Motor control
  - Memory and learning
  - Immunity and inflammatory responses
  - Neuroprotection




# CB1 receptors in the brain

Dense: Basal Ganglia (motor control), Cerebellum (coordination),  
Hippocampus (Memory and learning)

Middle Prefrontal and Parietal Cortex

Moderate: Amygdala (Anxiety), Spinal Cord (Pain modulation), Brainstem  
(Vomiting)



# CB1 receptors

- Typically the receptors exist on the axon terminal rather than on the postsynaptic cell
  - Affects many neurotransmitter systems
    - Acetylcholine
    - Dopamine
    - Norepinephrine
    - Serotonin
    - Glutamate
    - GABA

**The Human Endocannabinoid System**  
THC and CBN are known to "fit" like lock and key into network of existing receptors. The Endocannabinoid System exists to receive cannabinoids produced inside the body called "Anandamide" and "2-Arachidonyl-glycerol". Stimulating the ECS with plant-based cannabinoids restores balance and helps maintain symptoms.

CB1 receptors are concentrated in the brain and central nervous system but also sparsely populates other parts of the human body.

Receptors are found on cell surfaces

**THC**  
Tetrahydrocannabinol

**CB1**

**CBD**  
Cannabidiol

CBD does not directly "fit" CB1 or CB2 receptors but has powerful indirect effects still being studied.

**CBN**  
Cannabinol

**CB2**

CB2 receptors are mostly in the peripheral organs especially cells associated with the immune system.

[www.the-human-solution.com](http://www.the-human-solution.com)

**Presynaptic (sending neuron)**  
Cannabinoid Receptor  
Neurotransmitters

**Receptors**  
Lipid Precursors (fat cells)





## Role of CB2 receptors

---

- Located throughout the immune system and related organs, like the spleen, tonsils, and thymus gland
- Less dense in brain (more CB1)
- Modulate inflammatory responses to intestinal diseases such as Crohns and IBS
- Decreases inflammation in Arthritis and Alzheimer's





# Role of CB2 receptors and pain

---

- CB2 agonists (AM1241) inhibit nociception without producing CNS effects
  - The effects do not cross-over to morphine effects
- CB2 appears to modulate:
  - Acute pain
  - Chronic inflammatory pain
  - Post surgical pain
  - Cancer pain
  - Pain associated with nerve injury











# Role of CBD

- Low affinity for CB1 and CB2 receptors
- Acts as a antagonist on the CB1 and CB2 sites
- Extends duration of Anandamide and 2-AG
- Extend the duration of the effects of THC
  - Via inhibition of the cytochrome P450 enzyme system
- Dravet syndrome
- Other treatment-resistant epilepsies
- Pain
- Cancer
- Inflammation



# Endocannabinoid System: Effects of CB<sub>1</sub> Antagonism

	Site of Action	Mechanism(s)	Clinical Implications
	Hypothalamus/ nucleus accumbens	↓ Food intake	↓ Body weight ↓ Waist circumference
	Adipose tissue	↑ Adiponectin ↓ Lipogenesis	↓ Visceral fat ↓ Dyslipidemia ↓ Insulin resistance
	Muscle	↑ Glucose uptake ↑ O <sub>2</sub> consumption	↓ Insulin resistance
	Liver	↓ Lipogenesis	↓ Dyslipidemia ↓ Insulin resistance
	GI tract	↑ Satiety signals	↓ Body weight
	Pancreas	Potential effect on insulin secretion	Potential effect on glucose/insulin metabolism



# Summary

- The endocannabinoid system has been around million of years
- The use of cannabinoids may help chronic disease
- More research is now being conducted
- More to come in the future



# References

- Di Marzo, V., Melck, D., Bisogno, T., & De Petrocellis, L. (1998). Endocannabinoids: endogenous cannabinoid receptor ligands with neuromodulatory action. *Trends in neurosciences*, 21(12), 521-528.
- Ibsen, M. S., Connor, M., & Glass, M. (2017). Cannabinoid CB1 and CB2 Receptor Signaling and Bias. *Cannabis and Cannabinoid Research*, 2(1), 48-60.
- Mechoulam, R., Fride, E., & Di Marzo, V. (1998). Endocannabinoids. *European journal of pharmacology*, 359(1), 1-18.
- Zuardi, A. W. (2006). History of cannabis as a medicine: a review. *Revista Brasileira de Psiquiatria*, 28(2), 153-157.

